

Best practices in treating chronic noncancer pain

Since the mid-1990s, physicians have been increasingly prescribing higher doses and stronger opioids for their patients, particularly those with chronic noncancer pain.^{1,2} Opioid overprescription, improper risk assessment, and lack of monitoring opioid medication use have led to a significant health crisis in North America. Negative consequences, including opioid dependence, addiction, overdose, intentional or unintentional death, and diversion of drugs within the community, have impacts at both individual and societal levels. The Centers for Disease Control and Prevention reports that while death rates for conditions such as heart disease and cancer have decreased substantially over the past decade, the death rate associated with opioid pain prescription has increased significantly,³ despite the paucity of evidence to support the use of opioids to treat chronic noncancer pain.^{4,5} This increase in opioid pain prescriptions has likely contributed to the opioid crisis in which we find ourselves.

WorkSafeBC's practical experience supports the research findings. We seldom see long-term improvements in pain and function with long-term use of opioids. Instead, strong evidence supports the use of non-pharmacological options as first-line treatments for patients with chronic noncancer pain,⁶ which can be combined with pharmacological options or used on their own. The Centre for Effective Practice in Toronto divides nonpharmacological therapies into four categories:⁷

- Physical activity and exercise pro-

grams, including low-impact activities such as yoga, walking, and aquatic therapy.

- Physical therapies such as manual therapy.
- Psychological therapies such as cognitive behavioral therapy.

While death rates for conditions such as heart disease and cancer have decreased substantially over the past decade, the death rate associated with opioid pain prescription has increased significantly, despite the paucity of evidence to support the use of opioids to treat chronic noncancer pain.

- Self-management programs (treatment approaches that encourage individuals to be proactive in the care of their condition through lifestyle and behavioral changes and appropriate interaction with health care services).⁸

Non-opioid pharmacological treatments can also be effective, particularly when combined with some of these nonpharmacological options.

Finding the right treatment or combination of treatments requires careful trial, monitoring, and adjustment to ensure improvements in pain

and function are being achieved. While time consuming, this process is valuable to the patient.

WorkSafeBC's best practices seminars

Throughout 2018, WorkSafeBC will host community seminars to equip physicians and nurse practitioners with current best practices in the appropriate evidence-informed management of chronic noncancer pain. Our goal is to reduce harm and improve functional outcomes. Our informative, interactive sessions (Not Just a Prescription Pad: A Multimodal Approach to Chronic Pain Management) will give you practical knowledge and skills to apply to your practice. This workshop aligns with the College of Physicians and Surgeons of BC's standards and guidelines for safe prescribing of opioids, and physicians who attend will receive 2.5 Mainpro+ credits.

Sessions have already been held in Richmond, Burnaby, Kelowna, and Kamloops; the next session will be in Nelson on 18 June. Tentative dates for the remaining 2018 sessions are:

- 10 September: Langley/Surrey
- 25 September: Penticton/South Okanagan
- 2 October: Abbotsford/Chilliwack/Mission
- 19 October: Victoria
- 1 November: Courtenay/Comox/Campbell River
- 20 November: Delta/White Rock/Tsawwassen

All are evening sessions. Register online at <http://events.eply.com/chronicpain>, or call 1 877 231-8765. Seating is limited. Sessions are tentatively planned in 2019 for Coquitlam, Cranbrook/Fernie/Kimberley, Lillooet, Nanaimo/Qualicum, Port Moody,

Continued on page 269

This article is the opinion of WorkSafeBC and has not been peer reviewed by the BCMJ Editorial Board.

Continued from page 244

Prince George, Terrace, Vancouver, Vernon, and Williams Lake.

—**Peter Rothfels, MD**
Chief Medical Officer and Director
of Clinical Services, WorkSafeBC

References

1. Borgundvaag B, McLeod S, Khuu W, et al. Opioid prescribing and adverse events in opioid-naïve patients treated by emergency physicians versus family physicians: A population-based cohort study. *CMAJ Open* 2018;6:E110-E117.
2. Guy GP Jr, Zhang K, Bohm MK, et al. Vital signs: Changes in opioid prescribing in the United States, 2006–2015. *MMWR Morb Mortal Wkly Rep* 2017;66:697-704.
3. Rudd RA, Seth P, David F, Scholl L. Increases in drug and opioid-involved overdose deaths—United States, 2010–2015. *MMWR Morb Mortal Wkly Rep* 2016; 65:1445-1452.
4. McNicol ED, Midbari A, Eisenberg E. Opioids for neuropathic pain. *Cochrane Database Syst Rev* 2013;(8):CD006146.
5. Els C, Jackson TD, Hagtvedt R, et al. High-dose opioids for chronic noncancer pain: An overview of Cochrane Reviews. *Cochrane Database Syst Rev* 2017;10:CD012299.
6. Babatunde OO, Jordan JL, Van der Windt DA, et al. Effective treatment options for musculoskeletal pain in primary care: A systematic overview of current evidence. *PLoS One* 2017;12:e0178621.
7. Centre for Effective Practice. Management of chronic non-cancer pain tool. Toronto: Centre for Effective Practice, 2017. Accessed 4 April 2018. <https://thewellhealth.ca/cncp>.
8. Hernon MJ, Hall AM, O'Mahony JF, et al. Systematic review of costs and effects of self-management interventions for chronic musculoskeletal pain: Spotlight on analytic perspective and outcomes assessment. *Phys Ther* 2017;97:998-1019.

Continued from page 245

References

1. Polivy J, Herman CP. Mental health and eating behaviours: A bi-directional relation. *Can J Public Health* 2005;96 Suppl 3:S43-S46.
2. Li Y, Lv MR, Wei YJ, et al. Dietary patterns and depression risk: A meta-analysis. *Psychiatry Res* 2017;253:373-382.
3. Wallace CJK, Millev R. The effects of probiotics on depressive symptoms in humans: A systematic review. *Ann Gen Psychiatry* 2017;16:14.
4. Health Canada. About Natural Health Product Regulation in Canada. Accessed 5 March 2018. www.canada.ca/en/health-canada/services/drugs-health-products/natural-non-prescription/regulation.html.
4. Opie R, O'Neil A, Jacka FN, et al. A modified Mediterranean dietary intervention for adults with major depression: Dietary protocol and feasibility study from the SMILES trial. *Nutr Neurosci* 2017(April 19):1-15.

Continued from page 243

stool, which can be an early sign of colon cancer.

- Screening is only recommended for people who are not experiencing symptoms of colon cancer. Symptoms can include blood in the stool, abdominal pain, change in bowel habits, or unexplained weight loss. Anyone experiencing these symptoms should talk to their doctor about diagnostic testing they may need.
- Factors that put people at greater risk include having a first-degree relative (parent, sibling, or child) diagnosed under the age of 60, two or more first-degree relatives diagnosed at any age, and a personal history of adenomas.

For more information on the colon screening program, visit BC Cancer's screening website at www.screeningbc.ca.

**1 year. Unlimited
 number of trips.*
 1 low price.**



MEDOC® Travel Insurance gives you coverage for unlimited number of trips for an entire year, for about the same cost as insuring two separate trips!¹

1-855-734-8523
www.johnson.ca/doctorsofbc

JOHNSON 

Johnson Inc. ("Johnson") is a licensed insurance intermediary. MEDOC® is a Registered Trademark of Johnson. This insurance product is underwritten by Royal & Sun Alliance Insurance Company of Canada ("RSA") and administered by Johnson. The eligibility requirements, terms, conditions, limitations and exclusions, *(including but not limited to trip duration and intra-provincial trips), which apply to the described coverage are as set out in the policy. Policy wordings prevail. ¹Based on a comparison of MEDOC's multi-trip annual plan design against single trip plans with similar benefits. Johnson and RSA share common ownership. Call 1-800-563-0677 for details.

0266_17