

Supporting injured workers with PTSD

Most Canadians will be exposed to at least one traumatic event in their lifetime. Nevertheless, as the lifetime prevalence of posttraumatic stress disorder (PTSD) is reported to be 9.2% in Canada, the majority of persons exposed to trauma do not develop PTSD but rather have a normal response to an abnormal situation.¹⁻³ Yet the significant impact of repetitive trauma exposure on Canada's workforce clearly takes a toll. Rates of PTSD are greater in specific populations, such as first responders, than in the general population,¹ and compensable mental health injuries in BC are common. In 2019 and 2020, 641 and 555 injured workers, respectively, had claims accepted for PTSD by WorkSafeBC.

Implications for treating physicians

First-line treatment for PTSD is trauma-focused psychotherapy, which includes exposure-based and/or cognitive restructuring interventions such as prolonged exposure, cognitive processing therapy, eye movement desensitization and reprocessing, and trauma-focused cognitive-behavioral therapy (CBT).³⁻⁶ Although these therapies differ in their methods and protocols, each uses a cognitive or behavioral technique to assist the affected individual in processing the index trauma and to mitigate its illness-inducing appraisal. These interventions are available through WorkSafeBC and are provided by contracted mental health care providers (psychologists, clinical counselors, and/or occupational therapists). These providers are located throughout the province to assist injured workers as close to their home community as possible. Services are currently provided in person or virtually, and online CBT educational programs are also available.

This article is the opinion of WorkSafeBC and has not been peer reviewed by the BCMJ Editorial Board.

Referrals can be made to comprehensive outpatient and residential interdisciplinary programs when indicated. Expedited referrals to specific WorkSafeBC-contracted psychiatrists are also available.

Rates of PTSD are greater in specific populations, such as first responders, than in the general population.

Pharmacotherapy can also be helpful, particularly in injured workers who do not wish to engage in psychotherapy, in those with more severe presentations, or to target specific symptoms. To date, selective serotonin reuptake inhibitors (fluoxetine, sertraline, and paroxetine have the strongest evidence) and venlafaxine are the recommended first-line interventions.^{4,6} A host of other agents have been used as mono- or augmentation therapy when first-line pharmacotherapy is ineffective. The use of benzodiazepines merits particular comment. These are frequently prescribed for injured workers exposed to trauma, although most treatment guidelines either question their usefulness or recommend avoiding them altogether.⁷ In addition to general concerns about dependency and adverse side effects, more specific concerns revolve around their potential role in reducing the efficacy of exposure interventions (although a 2017 review⁷ has questioned the evidence for this finding) and increasing the severity of symptoms that can be present in patients with PTSD.^{4,8} As long-term use raises increasing concerns of dependency and loss of efficacy, WorkSafeBC normally limits financial coverage of benzodiazepines to a maximum of 2 weeks postinjury.

Other selective comments regarding pharmacotherapy for PTSD are as follows:

- Prazosin is frequently prescribed for PTSD-related nightmares with reported good effect, although one commonly used guideline suggests there is still insufficient evidence regarding its efficacy.⁴
- Cannabis or cannabis derivatives are not currently recommended due to the lack of evidence-established efficacy as well as potential adverse effects.⁴

Treating physicians are encouraged to consult recent clinical practice guidelines⁴⁻⁶ for further details and to follow the emerging literature related to cannabis and psychedelics for PTSD.^{4,9,10} Ultimately, treatment decisions need to be made on a case-by-case basis.

Finally, treating physicians may also wish to discuss posttraumatic growth—positive personal growth and resiliency after exposure to adversity—to normalize the experience and prevent pathology.^{11,12} This is not to minimize the significant impact of trauma exposure on workers who continue to provide invaluable service to our communities, particularly during recent global challenges.

For further assistance

Contact WorkSafeBC's RACE line/app Monday to Friday 8 a.m. to 5 p.m., billable under accepted or pending claims (billing code 19930). ■

—Tanya Fairweather, MD, CCFP, FCFP
Medical Advisor II

—Harry Karlinsky, MD, MSc, FRCRP
Psychiatric Consultant, WorkSafeBC

References

1. Carleton RN, Affi TO, Turner S, et al. Exposures to potentially traumatic events among public safety personnel in Canada. *Can J Behav Sci* 2019;51:37-52.
2. Duckers M, Alisic E, Brewin C. A vulnerability paradox in the cross-national prevalence of post-traumatic stress disorder. *Br J Psychiatry* 2016;209:300-305.
3. Shalev A, Liberzon I, Marmar C. Post-traumatic stress disorder. *N Engl J Med* 2017;376:2459-2469.

4. Department of Veterans Affairs, Department of Defense. VA/DoD clinical practice guideline for the management of posttraumatic stress disorder and acute stress disorder. June 2017. Accessed 15 October 2021. www.healthquality.va.gov/guidelines/MH/ptsd/VADoDPTSDCPGFinal012418.pdf
5. National Institute for Health and Care Excellence. Post-traumatic stress disorder. NICE Guideline. 5 December 2018. Accessed 15 October 2021. www.nice.org.uk/guidance/ng116.
6. Phoenix Australia. Australian guidelines for the prevention and treatment of acute stress disorder, posttraumatic stress disorder and complex PTSD. Accessed 15 October 2021. www.phoenixaustralia.org/australian-guidelines-for-ptsd.
7. Starcevic V. No role for benzodiazepines in post-traumatic stress disorder? A surplus of certainty despite scarce evidence. *Australas Psychiatry* 2017;25:339-341.
8. Guina J, Rossetter SR, DeRhodes BJ, et al. Benzodiazepines for PTSD: A systematic review and meta-analysis. *J Psychiatr Pract* 2015;21:281-303.
9. Tupper, Wood E, Yensen R, Johnson MW. Psychedelic medicine: A re-emerging therapeutic paradigm. *CMAJ* 2015;187:1054-1059.
10. Krediet, Bostoen T, Brecksemaet J, et al. Reviewing the potential of psychedelics for the treatment of PTSD. *Int J Neuropsychopharmacol* 2020;23:385-400.
11. Brooks S, Amlôt R, Rubin GJ, Greenberget N. Psychological resilience and post-traumatic growth in disaster-exposed organisations: Overview of the literature. *BMJ Mil Health* 2020;166:52-56.
12. Finstad GL, Giorgi G, Lulli LG, et al. Resilience, coping strategies and posttraumatic growth in the workplace following COVID-19: A narrative review on the positive aspects of trauma. *Int J Environ Res Public Health* 2021;18:9453.

Continued from page 433

- A dedicated space for physicians' children at smaller division meetings and supervised child-friendly activities/play stations at larger events hosted by the Thompson Region Division of Family Practice to help support member attendance and engagement.
- A new healthy snack program at Powell River General Hospital introduced by the Powell River Division of Family Practice after learning that some of its members felt unsafe to leave the facility to get meals while challenged with managing patient loads.

Doctors are encouraged to connect with their division, MSA, or RCCbc to learn more about supports for physician wellness in their area. For resources from the JCCs, visit www.collaborateonhealthbc.ca. ■

—Ahmer Karimuddin, MD

—Alan Ruddiman, MB BCH

References

1. Khan N, Palepu A, Dodek P, et al. Cross-sectional survey on physician burnout during the COVID-19 pandemic in Vancouver, Canada: The role of gender, ethnicity, and sexual orientation. *BMJ Open* 2021;11:e050380.
2. Walters L, Couper I, Stewart RA, et al. The impact of interpersonal relationships on rural doctors' clinical courage. *Rural Remote Health* 2021;21:6668.

Continued from page 432

References

1. McDonald HI, Tessier E, White JM, et al. Early impact of the coronavirus disease (COVID-19) pandemic and physical distancing measures on routine childhood vaccinations in England, January to April 2020. *Euro Surveill* 2020;25:pii=2000848.
2. Patel Murthy B, Zell E, Kirtland K, et al. Impact of the COVID-19 pandemic on administration of selected routine childhood and adolescent vaccinations - 10 U.S. jurisdictions, March–September 2020. *MMWR Morb Mortal Wkly Rep* 2021;70:840-845.
3. Causey K, Fullman N, Sorensen RJD, et al. Estimating global and regional disruptions to routine childhood vaccine coverage during the COVID-19 pandemic in 2020: A modelling study. *Lancet* 2021;398(10299):522-534.
4. World Health Organization. COVID-19 pandemic leads to major backsliding on childhood vaccinations, new WHO, UNICEF data shows. 15 July 2021. Accessed 8 November 2021. www.who.int/news/item/15-07-2021-covid-19-pandemic-leads-to-major-backsliding-on-childhood-vaccinations-new-who-unicef-data-shows.
5. BC Centre for Disease Control, BC Ministry of Health. Continuity, prioritization and safe delivery of immunization services during COVID-19 response. Updated 15 October 2021. Accessed 8 November 2021. www.bccdc.ca/resource-gallery/Documents/Guidelines%20and%20Forms/Guidelines%20and%20Manuals/Epid/CD%20Manual/Chapter%202%20-%20Imms/Continuity_of_Immunization_Services_During_COVID-19.pdf.

Available for streaming on all podcast platforms



BURNOUT AND COVID-19

Warning signs and when to act

with guests
Dr Jennifer Russel
and Dr Lawrence Yang

DocTalks
A Doctors of BC Podcast