

Myths versus facts: COVID-19 vaccine effects on pregnancy, fertility, and menstruation

Some of the most prevalent myths and the latest facts about fertility and COVID-19.

Jocelyn Y. Wong, Chelsea Elwood, MD, FRCSC, Deborah Money, MD, FRCSC, Caitlin Dunne, MD, FRCSC

Reproductive-age women and birthing persons may be hesitant to receive the COVID-19 vaccines due to concerns about hypothetical effects on pregnancy and fertility. Community physicians are well positioned to help share reliable and trustworthy

advice with patients who may have been exposed to disinformation online. However, keeping up with the latest myths and facts can be challenging in an era where both science and social media move rapidly. This article addresses some of the most prevalent myths and shares the latest facts about fertility and COVID-19.

Fact: The COVID vaccine does not cause infertility.

Disinformation about vaccines and miscarriage appears to have flourished in December 2020, when a blog post surfaced citing that a former employee of Pfizer believed that antibodies resulting from the COVID-19 vaccine could attack the placenta.¹ Subsequently, a myth propagated that antibodies that recognize the SARS-CoV-2 spike protein could cross-react with placental proteins called syncytins. Syncytin-1 protein acts in cell-to-cell adhesion, playing an essential role during placental attachment to the uterus during pregnancy; therefore, these claims provoked fear of placental damage with COVID-19 vaccination.² Since then, the original blog post has been taken down, and this myth has been debunked in several ways. Researchers have investigated the coronavirus's spike protein and compared it with placental syncytin-1, finding no significant similarities in their amino acid sequences.³ They also looked at serum from women with COVID-19 and did not detect reactions between these patients' antibodies and the syncytin-1 protein.³ Furthermore, if these myths were true about COVID-19 antibodies attacking placental syncytins, the pandemic

would have theoretically led to mass sterilization and miscarriage in women naturally infected with the virus, which has not occurred.

Despite the accumulating safety data, false rumors have continued to spread about the COVID-19 vaccines and infertility, generating feelings of hesitancy among pregnant people and people who plan to conceive. Recently, vaccine clinical trials have demonstrated that the rate of accidental pregnancies in vaccinated groups (29 of 39 848 individuals, 0.073%) was not significantly different from that in unvaccinated groups (28 of 39 845 individuals, 0.070%).¹ These findings suggest that fertility rates do not decrease with COVID-19 vaccination. A recent statement from the Society of Obstetricians and Gynaecologists of Canada confirmed: "There is absolutely no evidence, and no theoretic reason to suspect that the COVID-19 vaccine could impair male or female fertility. These rumors are unfounded and harmful."⁴

Research has also been conducted to address concerns regarding the impact of COVID-19 infection on fertility and assisted reproductive technology outcomes. A retrospective cohort study published in *The Lancet* followed 65 patients positive for SARS-CoV-2 antibodies and 195 matched patients negative for SARS-CoV-2 antibodies who underwent assisted reproductive technology treatments.⁵ Clinical pregnancy rates in the control group (54 clinical pregnancies out of 110 embryo transfer [ET] cycles, 49.1%) were not significantly different from rates in patients with mild or asymptomatic COVID-19 infections

Ms Wong is a medical student at the University of British Columbia in the class of 2024. Dr Elwood is a clinician-scientist in gynecologic reproductive infectious diseases. She is also a clinical associate professor in the Department of Obstetrics and Gynaecology at UBC. Dr Money is a clinician-scientist in gynecologic reproductive infectious diseases. She is a professor in the Department of Obstetrics and Gynaecology at UBC and the lead for CANCOVID-Preg (a Canada-wide surveillance program studying the outcome following COVID-19 for pregnant women and their infants), the lead for a pan-Canadian SARS-CoV-2 serosurveillance study using antenatal sera, and the lead for COVERED (a national study of the safety, effectiveness, and acceptability of the COVID-19 vaccines in pregnant and lactating persons). Dr Dunne is a reproductive endocrinology and infertility subspecialist. She is a clinical associate professor at UBC and co-director of the Pacific Centre for Reproductive Medicine.

This article has been peer reviewed.

(14 pregnancies out of 32 ET cycles, 43.8%).⁵ Overall, results showed no differences in ovarian reserve, ovarian response, biochemical pregnancy rate, clinical pregnancy rate, early miscarriage rate, and implantation rate between the case and control groups, demonstrating that COVID-19 infection has no impact on female fertility or assisted reproductive technology outcomes.⁵ Another observational study looked at 1347 pregnant women positive for SARS-CoV-2 infection in which 1273 (94.5%) had spontaneous pregnancies and 74 (5.5%) had IVF pregnancies. The distribution of asymptomatic and symptomatic patients was similar between the two conception groups (43.2% of IVF pregnancies had asymptomatic infections versus 51.5% of spontaneous pregnancies with asymptomatic infections; $P = .166$), and the analysis of clinical presentation (mild to moderate symptoms, pneumonia, complicated pneumonia/shock) between the two groups showed no significant differences.⁶ Overall, the COVID-19 symptomatology and symptom severity were the same in both the IVF and spontaneous conception groups.⁶ Furthermore, recent investigations show that receiving the SARS-CoV-2 mRNA vaccine has no effects on patients' performance in their subsequent IVF cycle, specifically no differences in patients' ovarian reserve or developing gametes and embryos.⁷ Thirty-six couples received two doses of vaccines and resumed their subsequent IVF treatment cycle 7 to 85 days after the second vaccine.⁷ Three pregnancies were recorded in 10 patients who underwent ET, indicating an acceptable pregnancy rate (30% per transfer).⁷ In January 2022, a study published in *Obstetrics & Gynecology* included 222 vaccinated and 983 unvaccinated patients who underwent ovarian hyperstimulation cycles.⁸ Results show no association between COVID-19 vaccination and fertilization rates.⁸ Moreover, 214 vaccinated and 733 unvaccinated patients who underwent frozen-thawed embryo transfer also showed a lack of significant association between COVID-19 vaccination and clinical pregnancy (adjusted odds ratio [aOR] 0.79; 95% CI, 0.54-1.16).⁸ Community physicians might find it helpful to be aware of these emerging findings regarding assisted reproductive technology, COVID-19, and COVID-19 vaccines to

address patient questions with accurate information and guide clinical decision-making.

Fact: The COVID-19 vaccine does not increase the risk of miscarriage or stillbirth.

Some young women and birthing persons are reluctant to be vaccinated because they fear it might affect a current or future pregnancy. This myth has become so pervasive that even some health care professionals have reported delaying vaccination. For example, members of the BC Nurses' Union were reported to

Current research shows that COVID-19 vaccines do not cause infertility or increase the risk of miscarriage or stillbirth.

have expressed concerns about the effects of COVID-19 vaccines on fertility and pregnancy, which contributed to many young nurses and other health care workers in BC remaining unvaccinated.⁹ Several studies have debunked these myths and confirmed that the risk of miscarriage after receiving the COVID-19 vaccine was not increased compared to the general population.¹⁰ Correspondence published in the *New England Journal of Medicine* found that out of 2456 participants enrolled in the Centers for Disease Control and Prevention's v-safe COVID-19 Vaccine Pregnancy Registry, the cumulative risk of spontaneous abortion from 6 to less than 20 weeks of gestation was 14.1% (95% CI, 12.1-16.1). This is consistent with the expected risk of spontaneous abortion of 11% to 22% in all recognized pregnancies.¹⁰ Additionally, BORN Ontario reports the stillbirth rate in individuals who received one or more COVID-19 vaccine doses before or during pregnancy to be 0.35%, which is similar to background stillbirth rates of 0.4% to 0.6% in Ontario.¹¹ The University of Washington conducted a prospective cohort study including 17 525 participants who were pregnant, lactating, or planning a pregnancy at the time of their first dose of the COVID-19 vaccine.¹² Results show that the odds of having several reactions

were statistically significantly decreased in pregnant individuals (i.e., fever after Pfizer dose 2: OR 0.44; 95% CI, 0.38-0.52; $P < .001$ and after Moderna dose 2: OR 0.48; 95% CI, 0.40-0.57; $P < .001$) compared with individuals who were not pregnant or lactating.¹² Obstetrical symptoms were reported in 346 of 7809 (4.4%) pregnant participants after the first vaccine dose and in 484 of 6444 (7.5%) pregnant participants after the second dose.¹² Overall, vaccine reactions and experiences were well tolerated and similar between pregnant or lactating individuals versus age-matched individuals who were neither pregnant nor lactating.¹² These findings were consistent with early data from the Canadian COVID-19 Vaccine Registry for Pregnant and Lactating Individuals (COVERED), showing no adverse events associated with the vaccines.¹² A large registry published in the *New England Journal of Medicine* reported that among 35 691 v-safe participants aged 16 to 54 who identified as pregnant, local and systemic reactions (injection-site pain, fatigue, headache, myalgias) were similar to patterns observed in nonpregnant women.¹³ Of the 3958 participants enrolled in the v-safe pregnancy registry, 827 individuals completed their pregnancies, with 86.1% live births, 12.6% spontaneous abortions, 0.1% stillbirths, and 1.2% with other outcomes. These findings were comparable to incidences published in current literature; therefore, results do not indicate any adverse pregnancy or neonatal outcomes in those who received mRNA COVID-19 vaccines.¹³ In summary, all the data at hand confirm that COVID-19 vaccines do not increase miscarriage or stillbirths in pregnant persons, and current literature suggests no safety concerns in those who are pregnant or lactating.

In contrast, a considerable amount of data show that outcomes are worse in pregnant individuals infected with SARS-CoV-2. COVID-19 is more likely to adversely affect pregnant women, leading to a 7% to 11% risk of being hospitalized for COVID-19-related morbidity, and a 1% to 4% risk of requiring intensive care.¹⁴ A Canada-wide study, CANCOVID-Preg, reported that pregnant people with COVID-19 infection have a stillbirth rate of 1.06%, which is twice that of the general population, though recent data suggest

the numbers are closer to population-level rates of 0.5%.¹⁵ Another complication is premature delivery, occurring in around 21.8% of pregnant individuals infected with COVID-19.¹⁶ Pre-term birth can result in extended stays in the neonatal intensive care unit (NICU) and may even lead to infant mortality. Research has also shown an association between certain risk factors, including age (≥ 35 years old), asthma, obesity, diabetes, hypertension, and heart disease, which increases the risk of severe morbidity from COVID-19 infections in pregnant persons.¹⁴ Given the abundance of data available, there are strong recommendations from the Society of Obstetricians and Gynaecologists of Canada and the National Advisory Committee on Immunization for pregnant, breastfeeding, and planning-to-become-pregnant individuals to receive the COVID-19 vaccine.^{14,17} Community physicians continue to take on the challenging and vital role of addressing the public's concerns about vaccine safety and encouraging vaccinations in pregnant people and people who plan to conceive.

Fact: The COVID-19 vaccine does not have enduring effects on menstrual periods.

News media have raised concerns regarding COVID-19 vaccines and their effects on menstruation.¹⁸ Until recently, information surrounding abnormal vaginal bleeding after vaccination has been anecdotal, and vaccine manufacturers did not record or report side effects related to menstruation.¹⁹ However, patient experiences may prompt questions about the vaccines and their effects on the menstrual cycle.¹⁹ In particular, vaccine hesitancy in the young population has been driven by false claims that COVID-19 vaccines and changes in menstrual cycles could negatively impact future pregnancy.²⁰ These concerns highlight the need for reproductive health outcomes to be included in postvaccine surveillance. Five institutions are now funded by the National Institutes of Health to conduct studies to examine this relationship between vaccination and irregular menstruation, and to address concerns that may be preventing women from receiving their COVID-19 vaccines.²¹

In January 2022, a study published in *Obstetrics & Gynecology* included 3959 individuals

Key points

- Current research shows that COVID-19 vaccines do not cause infertility or increase the risk of miscarriage or stillbirth.
- COVID-19 infection in pregnant women increases the risk of hospitalization, ICU admissions, premature delivery, and NICU admissions.
- Several guidelines, including those from the Society of Obstetricians and Gynaecologists of Canada and the National Advisory Committee on Immunization, strongly recommend COVID-19 vaccination for pregnant, breastfeeding, and planning-to-become-pregnant individuals.
- Recent studies are reporting minimal impact of COVID-19 vaccination on menstruation.
- Community physicians have a challenging and important role to play in keeping up with emerging vaccine myths versus scientific data to address the public's concerns about vaccine safety and to increase the public's trust in vaccines.

(2403 vaccinated) who prospectively tracked their cycles on a smartphone application. Overall, vaccination was associated with a less-than-one-day change in menstrual cycle length (first dose 0.71-day increase; 98.75% CI, 0.47-0.94; second dose 0.91-day increase; 98.75% CI, 0.63-1.19).²² Additionally, a retrospective study included 177 patients in a menstrual analysis and found that a quarter of patients with COVID-19 infections had influences on menstrual volume (20% had a significant decrease in menstrual volume and 5% had an increase in menstrual volume). Further analysis showed that 18% of patients had prolonged menstrual cycles, 3% had shortened cycles, and 7% showed cycle disorders. Follow-up determined that 84% of participants returned to their normal menstrual volume and 99% returned to their regular menstrual pattern after 1 to 2 months, suggesting that menstrual changes due to COVID-19 were temporary and quick to resolve.²³ Likewise, in the UK, most individuals who observed menstrual changes after vaccination also reported that their cycles returned to normal and changes were transient.²⁴ In total, 51 211 suspected menstrual cycle reactions were recorded in the UK, which is relatively low given that 74.1 million doses of COVID-19 vaccines had been administered by May 2022.²⁴ These studies, together, increase our confidence in the minimal impact of COVID-19 vaccines on menstruation. Current theories of changes in menstrual cycles focus on changes in the

immune system after receiving vaccinations.²⁵ In particular, shedding of the uterine lining during menstruation is an inflammatory response that involves the immune system. Therefore, vaccines that function by activating an immune response may temporarily change the normal course of menstruation. However, it is expected that if changes are noted, these changes would be short-term. To confirm these theories, controlled studies will need to be conducted, and researchers should put greater emphasis on including menstrual cycle tracking in future vaccine clinical trials. ■

Competing interests

Dr Dunne was a member of the *BCMJ* Editorial Board when this article was written, and is now the journal's editor, but did not participate in making the publication decision regarding this article.

References

1. Male V. Are COVID-19 vaccines safe in pregnancy? *Nat Rev Immunol* 2021;21:200-201.
2. Huang Q, Li J, Wang F, et al. Syncytin-1 modulates placental trophoblast cell proliferation by promoting G1/S transition. *Cell Signal* 2013;25:1027-1035.
3. Lu-Culligan A, Iwasaki A. False rumors about COVID-19 vaccines are scaring women. *The New York Times*. 26 January 2021. Accessed 28 October 2021. www.nytimes.com/2021/01/26/opinion/covid-vaccine-rumors.html.
4. The Society of Obstetricians and Gynaecologists of Canada. SOGC statement on COVID-19 vaccination and fertility. 2021. Accessed 5 April 2022. https://sogc.org/common/Uploaded%20files/Latest%20News/EN_SOGCStatement_COVID-19Vaccination-Fertility.pdf.

References continued on page 358

If monkeypox infection is confirmed

Advise the patient to continue infection control measures until all lesions have healed (i.e., the scabs have fallen off and re-epithelialization has occurred). Local public health will follow up with the patient for case and contact management. Treatment is largely supportive and focused on symptoms. ■

—Troy Grennan, MD, MSc, FRCPC

—Agatha Jassem, PhD, (D)ABMM, FCCM

—Mel Krajden, OBC, MD, FRCPC

—Mark Gilbert, MD, MHS, FRCPC

References

1. Thornhill JP, Barkati S, Walmsley S, et al. Monkeypox virus infection in humans across 16 countries—April–June 2022. *N Engl J Med* 2022;387:679–691.
2. World Health Organization. Joint ECDC-WHO Regional Office for Europe monkeypox surveillance bulletin. Accessed 25 August 2022. <https://monkeypoxreport.ecdc.europa.eu>.
3. BC Centre for Disease Control. Monkeypox—Information for health care providers. Accessed 25 August 2022. www.bccdc.ca/health-professionals/clinical-resources/monkeypox.



BC Medical Journal
@BCMJournal

[Follow](#)

The BC Medical Journal provides continuing medical education through scientific research, review articles, and updates on contemporary clinical practice. #MedEd

 Prescribing #nature to improve #health. #PaRx is Canada's first national, #EvidenceBased nature #prescription program, an initiative of the BC Parks Foundation driven by health care professionals. @bcparksfdn

Read the article: bcmj.org/news/prescribing-nature-improve-health



Follow us on Twitter for regular updates 

Continued from page 356

5. Wang M, Yang Q, Ren X, et al. Investigating the impact of asymptomatic or mild SARS-CoV-2 infection on female fertility and in vitro fertilization outcomes: A retrospective cohort study. *EClinicalMedicine* 2021; 38:101013.
6. Engels Calvo V, Cruz Melguizo S, Abascal-Saiz A, et al. Perinatal outcomes of pregnancies resulting from assisted reproduction technology in SARS-CoV-2-infected women: A prospective observational study. *Fertil Steril* 2021;116:731–740.
7. Orvieto R, Noach-Hirsh M, Segev-Zahav A, et al. Does mRNA SARS-CoV-2 vaccine influence patients' performance during IVF-ET cycle? *Repro Biol Endocrinol* 2021;19:69 (2021).
8. Aharon D, Lederman M, Ghofranian A, et al. In vitro fertilization and early pregnancy outcomes after coronavirus disease 2019 (COVID-19) vaccination. *Obstet Gynecol* 2022;139:490–497.
9. Fenton C. Fertility top of concerns for unvaccinated nurses, despite evidence showing vaccine safety. *CityNews*. 16 September 2021. Accessed 21 November 2021. <https://vancouver.citynews.ca/2021/09/16/bc-nurses-vaccine-fertility-concerns>.
10. Zauche LH, Wallace B, Smoots AN, et al. Receipt of mRNA Covid-19 vaccines and risk of spontaneous abortion. *N Engl J of Med* 2021;385:1533–1535.
11. BORN Ontario. Number of live births and stillbirths among infants born in Ontario, by COVID-19 vaccination status. Accessed 26 December 2021. www.bornontario.ca/en/news/number-of-live-and-stillbirths-among-infants-born-in-ontario-by-covid-19-vaccination-status.aspx.
12. Kachikis A, Englund JA, Singleton M, et al. Short-term reactions among pregnant and lactating individuals in the first wave of the COVID-19 vaccine rollout. *JAMA Netw Open* 2021;4:e2121310.
13. Shimabukuro TT, Kim SY, Myers TR, et al. Preliminary findings of mRNA Covid-19 vaccine safety in pregnant persons. *N Engl J Med* 2021;384:2273–2282.
14. Poliquin V, Castillo E, Boucoiran I, et al. SOGC statement on COVID-19 vaccination in pregnancy. *Society of Obstetricians and Gynaecologists of Canada*, 2020. Accessed 20 November 2021. https://sogc.org/common/Uploaded%20files/Latest%20News/SOGC_Statement_COVID-19_Vaccination_in_Pregnancy.pdf.
15. Money D. Canadian surveillance of COVID-19 in pregnancy: Epidemiology, maternal and infant outcomes: Maternal and infant outcomes (March 1, 2020 to March 31, 2021) from five Canadian provinces. Report No. 4. Reproductive Infectious Diseases Program, 2021. Accessed 5 April 2022. https://med-fom-ridprogram.sites.olt.ubc.ca/files/2021/06/CANCOVID_Preg-Report-4-ON-BC-QC-MB-AB_FINAL.pdf.
16. Khalil A, Kalafat E, Benlioglu C, et al. SARS-CoV-2 infection in pregnancy: A systematic review and meta-analysis of clinical features and pregnancy outcomes. *EClinicalMedicine* 2020;25:100446.
17. National Advisory Committee on Immunization. Vaccination and pregnancy: COVID-19. Accessed 26 December 2021. www.canada.ca/en/public-health/services/immunization-vaccines/vaccination-pregnancy-covid-19.html.
18. CBC News. What we know about COVID-19 vaccines in pregnant people. Accessed 20 November 2021. www.cbc.ca/player/play/1947977795768.
19. Lu-Culligan A, Epstein RH. No, we don't know if coronavirus vaccines affect periods. *The New York Times*. 20 April 2021. Accessed 31 October 2021. www.nytimes.com/2021/04/20/opinion/coronavirus-vaccines-menstruation-periods.html.
20. Male V. Menstrual changes after Covid-19 vaccination. *BMJ* 2021;374:n2211.
21. Eunice Kennedy Shriver National Institute of Child Health and Human Development. Item of interest: NIH funds studies to assess potential effects of COVID-19 vaccination on menstruation. 30 August 2021. Accessed 21 November 2021. www.nichd.nih.gov/newsroom/news/083021-COVID-19-vaccination-menstruation.
22. Edelman A, Boniface ER, Benhar E, et al. Association between menstrual cycle length and coronavirus disease 2019 (COVID-19) vaccination: A US cohort. *Obstet Gynecol* 2022;139:481–489.
23. Li K, Chen G, Hou H, et al. Analysis of sex hormones and menstruation in COVID-19 women of child-bearing age. *Reprod Biomed Online* 2021;42:260–267.
24. Medicines & Healthcare products Regulatory Agency. Coronavirus vaccine—Weekly summary of yellow card reporting. Accessed 25 May 2022. www.gov.uk/government/publications/coronavirus-covid-19-vaccine-adverse-reactions/coronavirus-vaccine-summary-of-yellow-card-reporting.
25. Garcia de Leon R, Richard J, Galea L. Menstrual irregularities and the COVID-19 vaccine. Accessed 4 November 2021. <https://womenshealthresearch.ubc.ca/blog/menstrual-irregularities-and-covid-19-vaccine>.