## Mental health and well-being among pandemic-era youth in British Columbia: Risk and protective factors

t the onset of the COVID-19 pandemic, health professionals hypothesized that a shadow pandemic of mental illness was imminent.1 Mental health issues among adolescents have been increasing steadily in Canada since 2011.2 Concerns regarding the impact on adolescent mental health3 of school closures and limiting in-person contact with peers were warranted. Research indicates that adolescents were among the populations whose mental health has suffered the most during COVID.4-8 Developmental resilience science provides insight for supporting youth mental well-being during this challenging time; more than 5 decades of evidence has demonstrated that positive experiences with families, with peers, in schools, and in communities act as protective and promotive factors for youth mental health.9

Data collected through the Youth Development Instrument (YDI) offer insight into pandemic-era adolescent mental health and well-being in BC.10 The YDI is an annual self-reported survey administered in BC secondary schools that asks youth ages 15 to 17 about their mental health and well-being. The YDI also measures risk and protective factors for mental health and well-being in individual, family, peer, school, and community domains.11 The YDI was piloted in the spring of 2020, and youth participation has increased each year since

This article is the opinion of the BC Centre for Disease Control and has not been peer reviewed by the BCMJ Editorial Board.

its inception. To date, YDI participation has included approximately 26 000 youth from 33 school districts and 28 independent schools across all five health regions in BC [Table 1]. Here we highlight key mental health and well-being challenges and opportunities elucidated by YDI research during the pandemic.

BC adolescent subjective mental well-being appears to have declined since the onset of the pandemic [Figure]. As well, in our latest wave of data collected from January to March 2023, 38% of youth screened positive for moderate to severe symptoms of depression, and 38% of youth screened positive for generalized anxiety disorder [Table 1]. YDI results suggest that some populations of youth in BC are experiencing poorer mental health and well-being than others. Analyses of 2022 YDI data indicated that youth with pre-existing mental health issues, those who report lower family affluence, and those living in rural areas of BC have poorer mental health and well-being compared with their

peers [Table 2]. LGBTQIA2S+ youth and those identifying as female or nonbinary reported poorer mental health and well-being than those who identify as heterosexual or male. Similar to 2022 data, preliminary analyses of 2023 YDI data identified youth with pre-existing mental health conditions as having the poorest mental health and well-being of YDI participants. These findings align with extant research that suggests these groups have been experiencing poorer mental health and well-being prior to and during the COVID pandemic.7 Thus, addressing social and structural inequities is paramount to population mental health.<sup>12</sup>

A motivating factor for creating the YDI was to identify malleable protective and promotive factors for youth mental health and well-being. In our analyses of 2021 YDI data, important correlates of mental health and well-being for BC adolescents included sleep, positive home experiences (e.g., supportive adults at home, positive communication with family), supportive peers, positive school experiences (e.g., school belonging,

TABLE 1. YDI mental health and well-being trends over 3 years (2021–2023).

Year	N	Depression (PHQ-8)	Anxiety (GAD-2)	High mental well-being	Six to seven positive childhood experiences
2021	2 295	40%	43%	27%	35%
2022	9 255	40%	41%	23%	37%
2023	14 596	38%	38%	20%	32%

Notes: Scores of 10 or greater on the Patient Health Questionnaire-8 (PHQ-8)18 indicate moderate to severe depression. Scores of 3 or greater on the Generalized Anxiety Disorder-2 (GAD-2)<sup>19</sup> indicate generalized anxiety disorder. High mental well-being constitutes youth who score at least 28 out of 35 on the Warwick-Edinburgh Mental Wellbeing Scale<sup>20</sup> (i.e., 80% or more of the scale's maximum value). YDI research indicates having six or more positive childhood experiences is a protective factor for depression and anxiety and a promotive factor for mental well-being.5

school safety), and social and emotional competencies, such as self-concept and optimism, after controlling for demographics such as racial identity, gender identity, sexual orientation, and family affluence.<sup>13</sup> YDI results for 2022 were similar: positive childhood experiences, such as having supportive families, peers, and teachers, were linked to significantly greater mental well-being and life satisfaction and significantly lower symptoms of depression and anxiety among youth.14 Moreover, we found that having six or more positive childhood experiences protected youth with adverse childhood experiences, such as neglect or abuse, against poor mental health and well-being. Thus, finding ways to increase positive experiences at home, among peers, at school, and in the community may serve as an important population-level strategy for promoting mental well-being and preventing mental illness among youth. Similar to extant research conducted among adults,15 we observed that positive childhood experiences are additive: the more of them that youth experience, the better their self-reported mental health and well-being on the YDI.14

Implementing evidence-based policies and programs that promote positive childhood experiences has been shown to lead to better mental health and well-being outcomes—for example, whole-school social




FIGURE. Subjective well-being among YDI participants (2021–2023).

and emotional learning programs that include student, teacher, parent/guardian, and community components.<sup>16</sup> Furthermore, addressing social determinants of health, such as poverty and discrimination, may lead to more positive childhood experiences and reduce the number of adverse childhood experiences, thus positively influencing mental health and well-being among youth.<sup>17</sup> Many of these initiatives are already underway in BC schools (e.g., the Mental Health in Schools Strategy, the sexual orientation and gender identity curriculum, the K-12 Anti-Racism Action Plan). Similar initiatives for supporting families and communities and in health care are necessary to promote mental health and well-being at a population level. ■

—Jacqueline Maloney, PhD Postdoctoral Fellow, Faculty of Health **Sciences, Simon Fraser University** 

—David Long, MD **Child and Youth Psychiatrist** Senior Data Analyst, Faculty of Health Sciences, SFU

—Jenna Whitehead, PhD **YDI Implementation Coordinator, Faculty** of Health Sciences, SFU

—Judy Wu, MPH **Doctoral Student, Faculty of Health** Sciences, SFU

—Julia Kaufmann, BSc Master of Public Health Student, Faculty of **Health Sciences, SFU** 

—Jillian Herring, MPH **Research Assistant, Faculty of Health** Sciences, SFU

-Rachel Correia, MPH **Research Assistant, Faculty of Health** Sciences, SFU

—Hasina Samji, PhD **Senior Scientist in Population Mental** Well-Being, BC Centre for Disease Control **Assistant Professor, Faculty of Health** Sciences, SFU

TABLE 2. Subpopulation analyses of 2022 YDI data.

Subpopulation	n	Depression (PHQ-8)	Anxiety (GAD-2)
Pre-existing mental health condition	2615	73%	72%
LGBTQIA2S+	2192	64%	61%
Heterosexual	7063	33%	34%
Female or nonbinary	4654	53%	56%
Male	4583	26%	35%
Low family affluence	1461	50%	49%
Medium or high family affluence	7358	38%	39%
Rural	1102	45%	46%
Urban	8099	39%	40%

n = the number of people from the total 2022 YDI sample (N = 9255) who identified as being in each subpopulation. The table includes the proportion of each population that screened positive for moderate to severe depression on the Patient Health Questionnaire-8 (PHQ-8) or for generalized anxiety disorder on the Generalized Anxiety Disorder-2 (GAD-2).

## References

1. Marques L, Bartuska AD, Cohen JN, Youn SJ. Three steps to flatten the mental health need curve amid the COVID-19 pandemic. Depress Anxiety 2020:37:405-406.

References continued on page 264

## Shifting access to apps—availability through the College Library

**♦** he CPS app from the Canadian Pharmacists Association and the Essential Evidence Plus app are no longer available from the Canadian Medical Association after closure of the CMA Library due to a change in the organization's focus. Another medical app, Epocrates, has exited the Canadian market. College registrants with Library access may be interested in the following alternatives.

This article is the opinion of the Library of the College of Physicians and Surgeons of BC and has not been peer reviewed by the BCMJ Editorial Board.

The CPS app, a Canadian source of drug and therapeutics information, is available via the College Library at www.cpsbc.ca/ registrants/library/drug-tools. The webbased version of the app and the drug interaction checker Lexi-Interact can also be accessed from that web page.

Alternatives to the decision-support apps Epocrates and Essential Evidence Plus include BMI Best Practice and DynaMed, which are available via the College Library's point-of-care web page at www.cpsbc .ca/registrants/library/point-care-tools. UpToDate is available through the Divisions of Family Practice, the Specialist Services Committee, and health authorities. The alternatives are not identical to Epocrates, but there are similarities, particularly for BMI Best Practice in terms of streamlined text.

Essential Evidence Plus contains decision support and content from the Cochrane Database of Systematic Reviews. Individual Cochrane reviews may be accessed via the Read by QxMD app, along with many other articles, or the entire database may be searched online at www.cpsbc .ca/registrants/library/databases.

Instructions for accessing all mobile apps through the College Library are on the Mobile Apps page: www.cpsbc.ca/registrants/ library/mobile-apps. ■

—Karen MacDonell **Director, Library Services** 

## **BCCDC** (Continued from page 263)

- 2. Wiens K, Bhattarai A, Pedram P, et al. A growing need for youth mental health services in Canada: Examining trends in youth mental health from 2011 to 2018. Epidemiol Psychiatr Sci 2020;29:e115.
- 3. Dove N, Wong D, Smolina K, et al. Impact of school closures on learning, and child and family wellbeing. BCMJ 2020;62:338.
- 4. Couturier J, Norris M. The shadow pandemic: Eating disorders, youth, and COVID-19. J Adolesc Health 2023;72:321-322.
- 5. Deng J, Zhou F, Hou W, et al. Prevalence of mental health symptoms in children and adolescents during the COVID-19 pandemic: A meta-analysis. Ann NY Acad Sci 2023;1520:53-73.
- 6. Madigan S, Racine N, Vaillancourt T, et al. Changes in depression and anxiety among children and adolescents from before to during the COVID-19 pandemic: A systematic review and meta-analysis. JAMA Pediatr 2023;177:567-581.
- Samji H, Wu J, Ladak A, et al. Review: Mental health impacts of the COVID-19 pandemic on children and youth—A systematic review. Child Adolesc Ment Health 2022;27:173-189.
- 8. Zandy M, El Kurdi S, Samji H, et al. Mental healthrelated healthcare service utilisation and psychotropic drug dispensation trends in British Columbia during COVID-19 pandemic: A population-based study. Gen Psychiatr 2023;36:e100941.

- Masten AS, Narayan AJ, Wright MO. Resilience processes in development: Multisystem integration emerging from four waves of research. In: Handbook of resilience in children. Goldstein S, Brooks RB, editors. Springer, Cham; 2023.
- 10. Gorfinkel LR, Snell G, Long D, et al. Access to mental health support, unmet need and preferences among adolescents during the first year of the COVID-19 pandemic. Health Promot Chronic Dis Prev Can 2023;43:182-190.
- 11. Samji H, Long D, Wu J, et al. The Youth Development Instrument: A population level survey of adolescent mental health and well-being indicators. Poster presented at the Banff International Conferences on Behavioural Science, Banff, AB, 20 March 2023.
- 12. Mansfield KL, Ukoumunne OC, Blakemore SJ, et al. Missing the context: The challenge of social inequities to school-based mental health interventions. JCPP Adv 2023;3:e12165.
- 13. Samji H, del Casal M, Kaufmann J, et al. Youth mental health and protective factors during the COVID-19 pandemic. Poster presented at the Banff International Conferences on Behavioural Science, Banff, AB, 20 March 2023.
- 14. Samji H, Long D, Correia R, et al. Positive childhood experiences as protective and promotive factors for mental health. Poster presented at the

- Frayme Learning Institute, Vancouver, BC, 22 Feb-
- 15. Bethell C, Jones J, Gombojav N, et al. Positive childhood experiences and adult mental and relational health in a statewide sample: Associations across adverse childhood experiences levels. JAMA Pediatr 2019;173:e193007.
- 16. Greenberg MT. Evidence for social and emotional learning in schools. Learning Policy Institute. 6 March 2023. Accessed 24 July 2023. https:// learningpolicyinstitute.org/product/evidence -social-emotional-learning-schools-report.
- 17. Gnanapragasam SN, Astill Wright L, Pemberton M, Bhugra D. Outside/inside: Social determinants of mental health. Ir J Psychol Med 2023;40:63-73.
- 18. Kroenke K, Strine TW, Spitzer RL, et al. The PHQ-8 as a measure of current depression in the general population. J Affect Disord 2009;114:163-173.
- 19. Staples LG, Dear BF, Gandy M, et al. Psychometric properties and clinical utility of brief measures of depression, anxiety, and general distress: The PHQ-2, GAD-2, and K-6. Gen Hosp Psychiatry 2019; 56:13-18.
- 20. Tennant R, Hiller L, Fishwick R, et al. The Warwick-Edinburgh Mental Well-being Scale (WEMWBS): Development and UK validation. Health Qual Life Outcomes 2007;5:63.